

p -adic l -functions and sums of powers

TAEKYUN KIM

*Jangjeon Research Institute for Mathematical Sciences & Physics, Ju-Kong Building
103-Dong 1001-ho, 544-4 Young-chang Ri Hapcheon-Up Hapcheon-Gun Kyungnam,
678-802, Korea*

e-mail: tkim64@hanmail.net or tkim@kongju.ac.kr

ABSTRACT. In this paper, we give an explicit p -adic expansion of

$$\sum_{\substack{j=1 \\ (j,p)=1}}^{np} \frac{(-1)^j}{j^r}$$

as a power series in n . The coefficients are values of p -adic l -function for Euler numbers.

1. Introduction

Let p be a fixed prime. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$, cf. [1], [3], [6], [10]. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = p^{-1}$. Kubota and Leopoldt proved the existence of meromorphic functions, $L_p(s, \chi)$, defined over the p -adic number field, that serve as p -adic equivalents of the Dirichlet L -series, cf. [8], [10]. These p -adic L -functions interpolate the values

$$L_p(1-n, \chi) = -\frac{1}{n}(1 - \chi_n(p)p^{n-1})B_{n, \chi_n}, \quad \text{for } n \in \mathbb{N} = \{1, 2, \dots\},$$

where $B_{n, \chi}$ denote the n th generalized Bernoulli numbers associated with the primitive Dirichlet character χ , and $\chi_n = \chi w^{-n}$, with w the *Teichmüller* character, see [2, 3, 5, 6, 17, 20]. In [14], L. C. Washington have proved the below interesting formula:

$$\sum_{\substack{j=1 \\ (j,p)=1}}^{np} \frac{1}{j^r} = - \sum_{k=1}^{\infty} \binom{-r}{k} (pn)^k L_p(r+k, w^{1-k-r}),$$

where $\binom{-r}{k}$ is binomial coefficient. In the recent many authors have studied q -extension of Euler numbers and Bernoulli numbers (see [1, 4, 5, 9, 12, 13]). These q -extensions seem to be valuable and worthwhile in the areas of mathematical physics and mathematics (see [1, 4, 6, 13, 14, 15, 17, 18, 19]). By using q -Volkenborn integration, Kim gave the interesting properties of q -Bernoulli and Euler polynomials

Key words and phrases: Euler zeta function, Euler numbers.

2000 Mathematics Subject Classification: 11S80, 11B68, 11M99.

[8, 9, 10, 11] and Ryoo-Kim-Agarwal have investigated the properties of the q -extension of Euler numbers and polynomials by using “Mathematica package”, see [14, 15]. The problems to find the sums of powers of consecutive q -integers were suggested. Kim and Schlosser treated the formulae for the sums of powers of consecutive q -integers [7, 11, 16] and these formulae were used to give the q -extension of Washington’s p -adic L -functions and sums of powers (see [6, 10, 20]). In [11, 14], we found the interesting formulae “alternating sums of powers of consecutive integers ” which are related to Euler numbers and polynomials. By using these alternating sums of powers of consecutive integers, we try to construct the p -adic l -functions and sums of powers for Euler numbers and polynomials, corresponding to Washington and Kim (see [10, 20]). The purpose of this paper is to give alternating p -adic harmonic series in terms of n and p -adic l -function for Euler numbers.

2. A note on l -series associated with Euler numbers and polynomials

We begin with well known Euler polynomials $E_n(x)$.

Definition 1. Euler polynomials are defined by

$$\frac{2}{e^t + 1} e^{xt} = \sum_{n=1}^{\infty} \frac{E_n(x)}{n!} t^n,$$

$E_n(x)$ are called n -th Euler polynomials. For $x = 0$, $E_n = E_n(0)$ are called Euler numbers. By the definition of Euler polynomials, we easily see that,

$$E_l(x) = \sum_{n=0}^l \binom{l}{n} E_n x^{l-n} \in \mathbb{C}[x].$$

From the generating function of Euler polynomials $F(t, x) = \frac{2}{e^t + 1} e^{xt}$, we derive

$$\begin{aligned} F(t, x) &= 2e^{xt} \sum_{l=0}^{\infty} (-1)^l e^{lt} \\ &= 2 \sum_{l=0}^{\infty} (-1)^l e^{(l+x)t}. \end{aligned} \tag{1}$$

For $k \in \mathbb{N}$, we note that

$$\begin{aligned} \left. \frac{d^k}{dt^k} F(t, x) \right|_{t=0} &= 2 \sum_{l=0}^{\infty} (-1)^l \left. \frac{d^k}{dt^k} e^{(l+x)t} \right|_{t=0} \\ &= 2 \sum_{l=0}^{\infty} (-1)^l (l+x)^k. \end{aligned} \tag{2}$$

Therefore we can define the Euler zeta function as follows:

Definition 2. For $s \in \mathbb{C}$, we define Euler zeta function as

$$\zeta_E(s) = 2 \sum_{l=0}^{\infty} \frac{(-1)^l}{(l+x)^s}. \quad (3)$$

By using Definition 2 and (2), we obtain the following:

Proposition 3. For $k \in \mathbb{N}$, we have

$$\zeta_E(-k, x) = E_k(x). \quad (4)$$

For $f(=\text{odd}) \in \mathbb{N}$,

$$\begin{aligned} \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} &= \frac{2}{e^t + 1} e^{xt} = 2 \sum_{l=0}^{\infty} (-1)^l e^{(l+x)t} = 2 \sum_{a=0}^{f-1} \sum_{l=0}^{\infty} (-1)^{a+lf} e^{(a+lf+x)t} \\ &= \sum_{a=1}^f (-1)^a 2 \sum_{n=0}^{\infty} (-1)^l e^{ft(l+\frac{x+a}{f})} = \sum_{a=1}^f (-1)^a \sum_{n=0}^{\infty} E_n\left(\frac{x+a}{f}\right) \frac{f^n t^n}{n!} \\ &= \sum_{n=0}^{\infty} f^n \sum_{a=1}^f (-1)^a E_n\left(\frac{x+a}{f}\right) \frac{t^n}{n!}. \end{aligned}$$

Thus we note that

$$E_n(x) = f^n \sum_{a=1}^f (-1)^a E_n\left(\frac{x+a}{f}\right), \quad (5)$$

where $f(=\text{odd}) \in \mathbb{N}$. This (5) is so called *Distribution for Euler polynomials*.

$$2 \sum_{l=0}^{n-1} (-1)^l l^m = (-1)^{n+1} \sum_{l=0}^{m-1} E_l n^{m-l} \binom{m}{l} + ((-1)^{n+1} + 1) E_m.$$

In particular, if n is even, then

$$2 \sum_{l=0}^{n-1} (-1)^l l^m = - \sum_{l=0}^{m-1} E_l n^{m-l}.$$

Let s be a complex variable and let $a, F(=\text{odd})$ be integers with $0 < a < F$.

$$\begin{aligned} H(s, a|F) &= \sum_{\substack{m \equiv a(F) \\ m > 0}} \frac{(-1)^m}{m^s} = \sum_{n=0}^{\infty} \frac{(-1)^{nF+a}}{(a+nF)^s} = (-1)^a \sum_{n=0}^{\infty} \frac{(-1)^n}{(a+nF)^s} \\ &= (-1)^a \sum_{n=0}^{\infty} \frac{(-1)^n}{F^s \left(\frac{a}{F} + n\right)^s} = \frac{(-1)^a F^{-s}}{2} 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n + \frac{a}{F}} \\ &= \frac{(-1)^a F^{-s}}{2} \zeta_E\left(s, \frac{a}{F}\right). \end{aligned} \quad (6)$$

Note that

$$H(-n, a|F) = (-1)^a \frac{F^n}{2} E_n \left(\frac{a}{F} \right). \quad (7)$$

Let χ be the primitive Dirichlet character with conductor $f(=\text{odd}) \in \mathbb{N}$.

$$\begin{aligned} F_\chi(t) &= 2 \sum_{n=0}^{\infty} e^{nt} \chi(n) (-1)^n \\ &= 2 \sum_{a=0}^{f-1} \sum_{n=0}^{\infty} e^{(a+nf)t} \chi(a+nf) (-1)^{a+nf} \\ &= \sum_{a=0}^{f-1} e^{at} \chi(a) (-1)^a 2 \sum_{n=0}^{\infty} (-1)^n e^{nft} \\ &= 2 \sum_{a=0}^{f-1} e^{at} \chi(a) (-1)^a \left(\frac{1}{e^{ft} + 1} \right) \\ &= \frac{2 \sum_{a=0}^{f-1} e^{at} \chi(a) (-1)^a}{e^{ft} + 1} \\ &= \sum_{n=0}^{\infty} E_{n, \chi} \frac{t^n}{n!}. \end{aligned} \quad (8)$$

Thus, we can define the below generalized Euler number attached to χ .

Definition 4. Let χ be the Dirichlet character with conductor $f(=\text{odd}) \in \mathbb{N}$. Then we define the generalized Euler numbers attached to χ as follows;

$$\frac{2 \sum_{a=0}^{f-1} e^{at} \chi(a) (-1)^a}{e^{ft} + 1} = \sum_{n=0}^{\infty} E_{n, \chi} \frac{t^n}{n!}.$$

$E_{n, \chi}$ will be called the n -th generalized Euler numbers attach to χ .

From the Definition 4, we derive the below formula:

$$\begin{aligned} \sum_{n=0}^{\infty} E_{n, \chi} \frac{t^n}{n!} &= \frac{2 \sum_{a=0}^{f-1} e^{at} \chi(a) (-1)^a}{e^{ft} + 1} = \sum_{a=0}^{f-1} \chi(a) (-1)^a \left(\frac{2}{e^{ft} + 1} e^{at} \right) \\ &= \sum_{a=0}^{f-1} \chi(a) (-1)^a \sum_{n=0}^{\infty} E_n \left(\frac{a}{f} \right) \frac{f^n t^n}{n!} = \sum_{n=0}^{\infty} f^n \sum_{a=0}^{f-1} \chi(a) (-1)^a E_n \left(\frac{a}{f} \right) \frac{t^n}{n!}. \end{aligned}$$

By comparing the coefficients on both sides, we easily see that

$$E_{n, \chi} = f^n \sum_{a=0}^{f-1} \chi(a) (-1)^a E_n \left(\frac{a}{f} \right). \quad (9)$$

Definition 5. For $s \in \mathbb{C}$, we define Dirichlet's l -function as follows:

$$l(s, \chi) = 2 \sum_{n=1}^{\infty} \frac{\chi(n)(-1)^n}{n^s}.$$

Note that

$$F_{\chi}(t) = 2 \sum_{n=1}^{\infty} e^{nt} \chi(n)(-1)^n = \frac{2 \sum_{a=0}^{f-1} (-1)^a \chi(a) e^{at}}{e^{ft} + 1} = \sum_{n=0}^{\infty} E_{n,\chi} \frac{t^n}{n!}.$$

For $k \in \mathbb{N}$,

$$\begin{aligned} E_{k,\chi} &= \left. \frac{d^k}{dt^k} F_{\chi}(t) \right|_{t=0} = 2 \sum_{n=1}^{\infty} \chi(n)(-1)^n \left. \frac{d^k}{dt^k} e^{nt} \right|_{t=0} \\ &= 2 \sum_{n=1}^{\infty} \chi(n)(-1)^n n^k. \end{aligned} \quad (10)$$

By Definition 5 and (10), we easily see that $l(-k, \chi) = E_{k,\chi}$, where $k \in \mathbb{N}$. Therefore we obtain the following:

Proposition 6. *Let k be the positive integer. Then we have*

$$l(-k, \chi) = E_{k,\chi}. \quad (11)$$

Let χ be the Dirichlet character with conductor f (=odd) $\in \mathbb{N}$. Then we note that

$$l(s, \chi) = 2 \sum_{a=1}^f \chi(a) H(s, a|f) \quad (12)$$

In Eq. (12), we give a value of $l(s, x)$ at negative integer:

$$\begin{aligned} l(-n, \chi) &= 2 \sum_{a=1}^f \chi(a) H(-n, a|f) \\ &= 2 \sum_{a=1}^f \chi(a)(-1)^a \frac{f^n}{2} E_n \left(\frac{a}{f} \right) \\ &= f^n \sum_{a=0}^{f-1} \chi(a)(-1)^a E_n \left(\frac{a}{f} \right) \\ &= E_{n,\chi}. \end{aligned}$$

The function $H(s, a|f)$ will be called partial zeta function which interpolates Euler polynomials at negative integers. The values of $l(s, \chi)$ at negative integers are algebraic, hence may be regarded as lying in an extension of $\mathbb{Q}_p$. We therefore

look for a p -adic function which agrees with $l(s, \chi)$ at negative integers in later.

3. A note on p -adic l -function

We define $\langle x \rangle = \frac{x}{w(x)}$, where $w(x)$ is the Teichmüller character. When F (=odd) is a multiple of p and $(a, p) = 1$, we define

$$H_p(s, a|F) = \frac{(-1)^a}{2} \langle a \rangle^{-s} \sum_{j=0}^{\infty} \binom{-s}{j} \left(\frac{F}{a} \right)^j E_j,$$

for $s \in \mathbb{Z}_p$.

It is easy to see that

$$\begin{aligned} H_p(-n, a|F) &= \frac{(-1)^a}{2} \langle a \rangle^n \sum_{j=0}^n \binom{n}{j} \left(\frac{F}{a} \right)^j E_j \\ &= \frac{(-1)^a}{2} F^n w^{-n}(a) \sum_{j=0}^n \binom{n}{j} \left(\frac{a}{F} \right)^{n-j} E_j \\ &= \frac{(-1)^a}{2} F^n w^{-n}(a) E_n \left(\frac{a}{F} \right) \\ &= w^{-n}(a) H(-n, a|F), \end{aligned}$$

for all positive integers. Now we consider p -adic interpolation function for Euler numbers as follows;

$$l_p(s, \chi) = 2 \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) H_p(s, a|F)$$

for $s \in \mathbb{Z}_p$.

Let n be natural number. Then we have

$$\begin{aligned} l_p(-n, \chi) &= 2 \sum_{\substack{n=1 \\ (n,p)=1}}^F \chi(a) H_p(-n, a|F) \\ &= E_{n, \chi w^{-n}} - p^n \chi w^{-n}(p) E_{n, \chi w^{-n}} \\ &= (1 - p^n \chi w^{-n}(p)) E_{n, \chi w^{-n}}. \end{aligned}$$

In fact, we have the formula

$$l_p(s, \chi) = \sum_{a=1}^F (-1)^a \langle a \rangle^{-s} \chi(a) \sum_{j=0}^{\infty} \binom{-s}{j} \left(\frac{F}{a} \right)^j E_j,$$

for $s \in \mathbb{Z}_p$.

This is a p -adic analytic function and has the following properties for $\chi = w^t$.

$$l_p(-n, w^t) = (1 - p^n)E_n, \quad \text{where } n \equiv t \pmod{p-1},$$

$$l_p(s, w^t) \in \mathbb{Z}_p \quad \text{for all } s \in \mathbb{Z}_p, \quad \text{when } t \equiv 0 \pmod{p-1}.$$

If $t \equiv 0 \pmod{p-1}$, then $l_p(s_1, w^t) \equiv l_p(s_2, w^t) \pmod{p}$ for all $s_1, s_2 \in \mathbb{Z}_p$,

$$l_p(k, w^t) \equiv l_p(k+p, w^t) \pmod{p}.$$

It is easy to see that

$$\frac{1}{r+k-1} \binom{-r}{k} \binom{1-r-k}{j} = \frac{-1}{j+k} \binom{-r}{k+j-1} \binom{k+j}{j},$$

for all positive integers with r, j, k with $j, k \geq 0$, $j+k > 0$ and $r \neq 1-k$.

Thus, we note that

$$\frac{1}{r+k-1} \binom{-r}{k} \binom{1-r-k}{j} = \frac{1}{r-1} \binom{-r+1}{k+j} \binom{k+j}{j}.$$

Hence, we have

$$\frac{r}{r+k} \binom{-r+1}{k} \binom{-r-k}{j} = \binom{-r}{k+j} \binom{k+j}{j},$$

where k, j are positive integers. Let $F(=\text{odd})$ be positive integers. Then

$$\begin{aligned} & \sum_{l=0}^{n-1} \frac{(-1)^{Fl+a}}{(Fl+a)^r} \\ &= \sum_{l=0}^{n-1} (-1)^{Fl+a} a^{-r} \sum_{s=0}^{\infty} \binom{-r}{s} \left(\frac{Fl}{a}\right)^s \\ &= \sum_{m=0}^{\infty} \binom{-r}{m} a^{-r} \left(\frac{F}{a}\right)^m (-1)^a \sum_{l=0}^{n-1} (-1)^l l^m \\ &= \sum_{m=0}^{\infty} \binom{-r}{m} a^{-r} \left(\frac{F}{a}\right)^m (-1)^a \frac{(-1)^{n+1}}{2} \sum_{l=0}^{m-1} E_l n^{m-l} \binom{m}{l} + ((-1)^{m-1} + 1) E_m, \end{aligned}$$

when n is even integer

$$\begin{aligned}
& \sum_{l=0}^{n-1} \frac{(-1)^{Fl+a}}{(Fl+a)^r} \\
&= a^{-r} (-1)^a \sum_{m=0}^{\infty} \binom{-r}{m} \left(\frac{F}{a}\right)^m \left\{ \frac{(-1)^{m+1}}{2} \sum_{l=0}^{m+1} E_l n^{m-l} \binom{m}{l} \right\} \\
&= -a^{-r} (-1)^a \sum_{s=0}^{\infty} \binom{-r}{s} \left(\frac{F}{a}\right)^s \left(\sum_{l=0}^{s+1} E_l n^{s-l} \binom{s}{l} \right) \\
&= - \sum_{s=0}^{\infty} \binom{-r}{s} w^{-r}(a) \left(\frac{F}{a}\right)^s \frac{(-1)^a}{2} \langle a \rangle^{-r} \sum_{l=0}^{s-1} E_l n^{s-l} \binom{s}{l} \\
&= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \binom{-r}{k+l} w^{-r}(a) \left(\frac{a}{F}\right)^{-k-l} n^{k+l} \frac{(-1)^a}{2} \langle a \rangle^{-r} E_l \left(\frac{1}{n}\right)^l \binom{k+l}{l} \\
&= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{r}{r+k} \binom{-r-1}{k} \binom{-r-k}{l} a^{-r} \left(\frac{a}{F}\right)^{-k-l} n^{k+l} \frac{(-1)^a}{2} E_l \left(\frac{1}{n}\right)^l \\
&= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{r}{r+k} \binom{-r-1}{k} w^{-k-r}(a) (nF)^k \frac{(-1)^a}{2} \langle a \rangle^{-k-r} \sum_{l=0}^{\infty} \binom{-r-k}{l} E_l \left(\frac{F}{a}\right)^l \\
&= - \sum_{k=0}^{\infty} \frac{r}{r+k} \binom{-r-1}{k} w^{-k-r}(a) (nF)^k H_p(r+k, a|F).
\end{aligned}$$

For $F = p$, $r, n(=\text{even}) \in \mathbb{N}$, we see that

$$\begin{aligned}
2 \sum_{\substack{j=1 \\ (j,p)=1}}^{np} \frac{(-1)^j}{j^p} &= 2 \sum_{a=1}^{p-1} \sum_{l=0}^{n-1} \frac{(-1)^{a+pl}}{(a+pl)^r} \\
&= -2 \sum_{a=1}^{p-1} \sum_{k=0}^{\infty} \frac{r}{r+k} \binom{-r-1}{k} w^{-k-r}(a) (pn)^k H_p(r+k, a|p) \\
&= - \sum_{k=0}^r \frac{r}{r+k} \binom{-r-1}{k} (pn)^k 2 \sum_{a=1}^{p-1} w^{-k-r}(a) H_p(r+k, a|p) \\
&= - \sum_{k=0}^r \binom{-r-1}{k} (pn)^k l_p(r+k, w^{-k-r}) \\
&= - \sum_{k=0}^{\infty} \frac{r}{r+k} \binom{-r-1}{k} (pn)^k l_p(r+k, w^{-k-r}).
\end{aligned}$$

Therefore we obtain the following:

Theorem 6. *Let p be an odd prime and let $n(=even)$, $r \in \mathbb{N}$. Then we have*

$$2 \sum_{\substack{j=1 \\ (j,p)=1}}^{np} \frac{(-1)^j}{j^r} = - \sum_{k=0}^{\infty} \frac{r}{r+k} \binom{-r-1}{k} (pn)^k l_p(r+k, w^{-k-r}).$$

References

- [1] L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J., **15**(1948), 987-1000.
- [2] J. Diamond, *The p-adic log gamma function and p-adic Euler constant*, Trans. Amer. Math. Soc., **233**(1977), 321-337.
- [3] B. Ferrero and R. Greenberg, *On the behavior of p-adic L-functions at $s = 0$* , Invent. Math., **50**(1978), 91-102.
- [4] A.S. Hegazi, M. Mansour, *A Note on q-Bernoulli Numbers and Polynomials*, Journal of Nonlinear Mathematical Physics, **13**(2006), 9-18.
- [5] K. Iwasawa, *Lecture notes on p-adic L-function*, Princeton Univ. Press (1972).
- [6] T. Kim, *On explicit formulas of p-adic q-L-functions*, Kyushu J. Math., **48**(1994), 73-86.
- [7] T. Kim, *Sums powers of consecutive q-integers*, Advan. Stud. Contemp. Math., **9**(2004), 15-18.
- [8] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys., **9**(2002), 288-299.
- [9] T. Kim, *Power series and Asymptotic series associated with the q-analogue of two-variable p-adic L-function*, Russian Journal of Mathematical Physics, **12**(2005), 186-195.
- [10] T. Kim, *On p-adic q-L-functions and sums of powers*, Discrete Math., **252**(2002), 179-187.
- [11] T. Kim, *A note on the alternating sums of powers of consecutive integers*, arXiv. Math. NT/0508233, **1**(2005), 1-4.
- [12] N. Koblitz, *A new proof of certain formulas for p-adic L-function*, Duke Math. J., **46**(1979), 455-468.
- [13] B. A. Kupersmidt, *Reflection symmetries of q-Bernoulli polynomials*, J. Nonlinear Math. Phys., **12**(1979), 412-422.
- [14] C.S. Ryoo, T. Kim, R.P. Agarwal, *A numerical investigation of the roots of q-polynomials*, International Journal of Computer Mathematics, **83**(2006), 223-234.
- [15] C.S. Ryoo, T. Kim, R.P. Agarwal, *Exploring the multiple Changhee q-Bernoulli polynomials*, International Journal of Computer Mathematics, **82**(2005), 223-234.
- [16] M. Schlosser, *q-analogues of the sums of consecutive integers, squares, cubes, quarts, and quints*, The Electronic J. Combinatorics, **11**(2004), R 71.

- [17] K. Shiratani and S. Yamamoto, *On a p -adic interpolation function for Euler numbers and its derivatives*, Mem. Fac. Sci, Kyushu Univ., **39**(1985), 113-125.
- [18] Y. Simsek, D. Kim and S.-H. Rim, *On the two-variable Dirichlet q - L -series*, Adv. Stud. Contemp. Math. (Kyungshang), **10(2)**(2005), 131-142
- [19] H. M. Srivastava, T. Kim, Y. Simsek, *q -Bernoulli numbers and polynomials associated with multiple q -zeta functions and basic L -series*, Russian Journal of Mathematical Physics, **12**(2005), 241-278
- [20] L. C. Washington, *p -adic L -functions and sums of powers*, J. Number Theory, **69(1)**(1988), 50-61.